



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

NOTE ON "AN APPLICATION OF ELLIPTIC FUNCTIONS TO GEOMETRY." *

BY PROF. JAS. H. BOYD, St. Paul, Minn.

Formulas (6) and (11), ANNALS OF MATHEMATICS, Vol. VI, pp. 95 and 97,
give

$$\begin{aligned} S_A(z) &= \sum_{+\infty}^{-\infty} n \frac{\pi \tan^4 \alpha \cdot h^{4n} z^4}{(h^{2n} z^2 - z_1^2)^2 (h^{2n} z^2 - z_2^2)^2} \\ &= C_1 p(u - u_1) + C_2 p(u - u_2) + C_3 \frac{\sigma'}{\sigma}(u - u_1) \\ &\quad + C_4 \frac{\sigma'}{\sigma}(u - u_2) + C_5, \end{aligned} \tag{1}$$

where $S_A(z)$ represents the sum of the areas of circles which touch each other successively and are inscribed in a crescent formed by the intersection of two circles. The C 's are constants to be determined.

We know by the theory of decomposition of fractions into their partial fractions, that,

$$\frac{x}{(x-a)^2(x-\beta)^2} = \frac{A_0}{(x-a)^2} + \frac{A_1}{x-a} + \frac{B_0}{(x-\beta)^2} + \frac{B_1}{x-\beta},$$

where,

$$\begin{aligned} A_0 &= \frac{a}{(a-\beta)^2}, & A_1 &= -\frac{a+\beta}{(a-\beta)^3}, \\ B_0 &= \frac{\beta}{(a-\beta)^2}, & B_1 &= \frac{a+\beta}{(a-\beta)^3}. \end{aligned} \tag{2}$$

Putting, therefore, $x = h^{2n} z^2$, $a = z_1^2$, and $\beta = z_2^2$, we shall have

$$\begin{aligned} \frac{h^{4n} z^4}{(h^{2n} z^2 - z_1^2)^2 (h^{2n} z^2 - z_2^2)^2} &= \frac{z_1^2 h^{2n} z^2}{(z_1^2 - z_2^2)^2 (h^{2n} z^2 - z_1^2)^2} \\ &\quad + \frac{z_2^2 h^{2n} z^2}{(z_1^2 - z_2^2)^2 (h^{2n} z^2 - z_2^2)^2} - \frac{(z_1^2 + z_2^2) h^{2n} z^2}{(z_1^2 - z_2^2)^3 (h^{2n} z^2 - z_1^2)} \\ &\quad + \frac{(z_1^2 + z_2^2) h^{2n} z^2}{(z_1^2 - z_2^2)^3 (h^{2n} z^2 - z_2^2)}; \end{aligned} \tag{3}$$

* See ANNALS OF MATHEMATICS, Vol. VI, No. 4.

hence substituting in (1),

$$\begin{aligned}
 S_A(z) &= C_1 p(u - u_1) + C_2 p(u - u_2) + C_3 \frac{\sigma'}{\sigma}(u - u_1) + C_4 \frac{\sigma'}{\sigma}(u - u_2) + C_5 \\
 &= \sum_{n=-\infty}^{+\infty} \frac{\pi \tan^4 \alpha \cdot z_1^2 h^{2n} z^2}{(z_1^2 - z_2^2)^2 (h^{2n} z^2 - z_1^2)^2} + \sum_{n=-\infty}^{+\infty} \frac{\pi \tan^4 \alpha \cdot z_2^2 h^{2n} z^2}{(z_1^2 - z_2^2)^2 (h^{2n} z^2 - z_2^2)^2} \\
 &\quad + \sum_{n=-\infty}^{+\infty} \frac{\pi \tan^4 \alpha \cdot (z_2^4 - z_1^4) \cdot h^{2n} z^2}{(z_1^2 - z_2^2)^3 (h^{2n} z^2 - z_1^2)(h^{2n} z^2 - z_2^2)}. \quad (4)
 \end{aligned}$$

From § 9, (4) of Weierstrass and Schwarz's Elliptic Function Formulas (Edition of 1885), we have

$$p u = -\frac{\eta}{\omega} - \left[\frac{\pi}{\omega} \right]^2 \left\{ \frac{z^2}{(z^2 - 1)^2} + \sum_n \frac{h^{2n} z^2}{(1 - h^{2n} z^2)^2} + \sum_n \frac{h^{2n} z^{-2}}{(1 - h^{2n} z^{-2})^2} \right\}. \quad (5)$$

We had found (ANNALS, Vol. VI, p. 96) that, when u becomes $u - u_1$ or $u - u_2$, z becomes $\frac{z}{z_1}$ or $\frac{z}{z_2}$ respectively.

Hence,

$$\begin{aligned}
 p(u - u_1) &= -\frac{\eta}{\omega} - \left[\frac{\pi}{\omega} \right]^2 \left\{ \frac{z^2 z_1^2}{(z^2 - z_1^2)^2} + \sum_n \frac{h^{2n} z^2 z_1^2}{(z_1^2 - h^{2n} z^2)^2} \right. \\
 &\quad \left. + \sum_n \frac{h^{2n} z^{-2} z_1^{-2}}{(z_1^{-2} - h^{2n} z^{-2})^2} \right\}, \quad (6)
 \end{aligned}$$

$$= -\frac{\eta}{\omega} - \left[\frac{\pi}{\omega} \right]^2 \cdot z_1^2 \cdot \sum_{n=-\infty}^{+\infty} \frac{h^{2n} z^2}{(h^{2n} z^2 - z_1^2)^2}, \quad (7)$$

$$p(u - u_2) = -\frac{\eta}{\omega} - \left[\frac{\pi}{\omega} \right]^2 \cdot z_2^2 \cdot \sum_{n=-\infty}^{+\infty} \frac{h^{2n} z^2}{(h^{2n} z^2 - z_2^2)^2}. \quad (8)$$

From page 96, Volume VI of the ANNALS, just below formula (9), we have

$$\begin{aligned}
 \sum_{n=-\infty}^{+\infty} \frac{h^{2n} z^2}{(h^{2n} z^2 - z_1^2)(h^{2n} z^2 - z_2^2)} &= \frac{\omega}{\pi i (z_1^2 - z_2^2)} \cdot \left\{ \frac{\sigma'}{\sigma}(u - u_1) - \frac{\sigma'}{\sigma}(u - u_2) \right\} \\
 &\quad + \frac{\eta}{\pi i} \left(\frac{u_1 - u_2}{z_1^2 - z_2^2} \right); \quad (9)
 \end{aligned}$$

and from (7) and (8),

$$\sum_{n=-\infty}^{+\infty} \frac{h^{2n} z^2}{(h^{2n} z^2 - z_1^2)^2} = - \left[\frac{\omega}{\pi z_1} \right]^2 \cdot p(u - u_1) - \frac{\omega \eta}{(\pi z_1)^2}, \quad (10)$$

$$\sum_{n=-\infty}^{+\infty} \frac{h^{2n} z^2}{(h^{2n} z^2 - z_2^2)^2} = - \left[\frac{\omega}{\pi z_2} \right]^2 \cdot p(u - u_2) - \frac{\omega \eta}{(\pi z_2)^2}. \quad (11)$$

Substituting the values of the \sum 's in (9), (10), and (11) in (4), we have,

$$\begin{aligned} & C_1 p(u - u_1) + C_2 p(u - u_2) + C_3 \frac{\sigma'}{\sigma}(u - u_1) + C_4 \frac{\sigma'}{\sigma}(u - u_2) + C_5 \\ &= - \frac{\omega^2}{\pi} \cdot \frac{\tan^4 \alpha}{(z_1^2 - z_2^2)^2} \cdot p(u - u_1) - \frac{\omega}{\pi} \cdot \frac{\eta \tan^4 \alpha}{(z_1^2 - z_2^2)^2} \\ & \quad - \frac{\omega^2}{\pi} \cdot \frac{\tan^4 \alpha}{(z_1^2 - z_2^2)^2} \cdot p(u - u_2) - \frac{\omega}{\pi} \cdot \frac{\eta \tan^4 \alpha}{(z_1^2 - z_2^2)^2} \\ & \quad - \frac{\omega \tan^4 \alpha}{i} \cdot \frac{z_1^2 + z_2^2}{(z_1^2 - z_2^2)^3} \left\{ \frac{\sigma'}{\sigma}(u - u_1) - \frac{\sigma'}{\sigma}(u - u_2) \right\} \\ & \quad - \frac{(z_1^2 + z_2^2)}{(z_1^2 - z_2^2)^3} \cdot \frac{\eta}{i} \cdot \frac{\tan^4 \alpha}{(u_1 - u_2)^{-1}}. \end{aligned} \quad (12)$$

The C 's, therefore, must have the following values :

$$C_1 = C_2 = - \frac{\omega^2}{\pi} \cdot \frac{\tan^4 \alpha}{(z_1^2 - z_2^2)^2}, \quad (13)$$

$$C_3 = - C_4 = - \frac{\omega \tan^4 \alpha}{i} \cdot \frac{z_1^2 + z_2^2}{(z_1^2 - z_2^2)^3}, \quad (14)$$

$$C_5 = \frac{\eta \tan^4 \alpha}{(z_1^2 - z_2^2)^2} \left\{ - \frac{2\omega}{\pi} - \frac{z_1^2 + z_2^2}{z_1^3 - z_2^2} \cdot \frac{u_1 - u_2}{i} \right\} \quad (15)$$

Our original problem is now completely solved, since $S_A(z)$ has been shown to depend upon p -function of $u - u_1$, $u - u_2$, and $\frac{\sigma'}{\sigma}$ -function of the same, and upon the coefficients C_1 , C_2 , C_3 , C_4 , C_5 , which involve z_1 , z_2 , u_1 , u_2 , and η , ω , α . How these latter elements may be calculated has already been shown in the ANNALS, Vol. VI, p. 97.